

Int. J. Forest, Soil and Erosion, 2016 6 (4)

ISSN 2251-6387

© February 2016, GHB's Journals, IJFSE, Shabestar, Iran

Research Paper

MEASURES OF BIODIVERSITY INDEX OF ALGAL COMMUNITIES IN SHADAWANKA RIVER BAUCHI - NIGERIA

Sindama, A

Department of Biological Sciences, College of Pure and Applied Sciences, Federal University, Wukari, Taraba State, Nigeria

E-mail: sindamaalex@yahoo.com

Abstract

The study of the diversity, species richness, and abundance of algae was undertaken for the period of eight months (February – September 2007) in Shadawanka River, Bauchi, Bauchi State. Assessment of physicochemical characteristics of the river was also attempted to find out the state of the water quality of the river. Four hundred and eighty one species of phytoplankton were recorded. This consisted of 276 species of diatoms which was made up of 237 species of Pennales and 39 species of Centrales. 123 species of desmids (green algae), 41 species of Euglenophyceae, 34 species of Cyanophyceae, 3 species of Phaeophyceae and Xanthophyceae having 4 species from a single order Heterococcales. A total of 201 algal species in the Epiphyton were identified, 122 species belonged to the Bacillariophyceae. This consisted of 97 species of Pennales and 25 species of Centrales. There were 58 species of Chlorophyceae, 14 species of Cyanophyceae, 7 species of Euglenophyceae and 3 species of Xanthophyceae respectively. Seventy one species of Epilithon were observed, which included 33 species of Bacillariophyceae consisted of 22 species of Pennales and 11 species of Centrales, 23 species of Chlorophyceae and 9 species of Cyanophyceae and 6 species of Euglenophyceae. The highest species richness (16.50) was obtained in the phytoplankton and the lowest (3.82) was in the Epilithon whilst Epiphyton was an average (10.51). The similarities between phytoplankton and Epiphyton and that between Epiphyton and Epilithon were same (0.17), whilst the similarities obtained between the phytoplankton and Epilithon communities was lower (0.14). The River is well oxygenated ($DO > 9.8 \text{ mg/L}$) and is Characterised by moderate nutrient levels, high electrical conductivity ($> 204 \mu\text{mScm}^{-1}$) and low BOD ($2.1\text{-}6.4 \text{ mg/L}$). pH was circumneutral, having three regimes, this attest that the water is not polluted and can support other forms of aquatic life

Keywords: Centrales, Physicochemical, Phytoplankton, Epilithon, Epiphyton.

Introduction

The preference for habitat characteristic by certain species of algae has been stressed by Patrick (1988) who claimed that the restriction of certain taxa to a given habitat might probably be as a result of a reflection of the peculiarity of different habitats. The Englenales were found to be restricted to confluences and downstream areas, possibly as a result of nutrient levels coupled with favourable temperatures, while desmids occupy a wide range of ecological niches (Moss, 1973). Studies have confirmed that, a change in species diversity and phytoplankton taxonomic composition is attributable to changes in nutrient levels (Watson *et al.*, 2007). Reynolds (1984) reported that, such changes are related to the differences between the taxa in terms of storage, nutrients uptake, growth and loss. Algae and other aquatic organisms are usually considered to be opportunistic species as they respond quickly to minor perturbations in the physical conditions of their environment (Nwankwo, 1988).

Materials and Methods

Shadawanka River is situated in the Shadawanka military barracks, 2km, off Jos road, east of Wuntin dada village, south of Shadawanka village. It resulted from two confluence rivers namely: Lafiyari and Zamfara rivers. Shadawanka River serves majorly as a source of water for livestock watering and irrigation purposes for both the dry and wet seasons. Aquatic macrophytes common in the river are; *Nymphia lotus L.*, *Pennesetum sp.*, *Andropogon sp.*, and *heteropogon sp.* The dry season usually ranges between the months of November to March while the Wet season last from April to October.

Collection of Samples

Water samples for physicochemical analysis were collected monthly for a period of eight months (February- September 2007) between 7 and 8 hours from four stations namely: Stations A, B, C and D (Figure 1) in 1 litre acid rinsed polyethene bottles and transferred to the laboratory in a cooler packed with ice prior to analysis. Samples for biological analysis was obtained as thus: Phytoplankton were collected using a plankton net of mesh size 55µm towed at low speed for 10min and immediately collected in screw capped bottles and immediately fixed with 4% formalin. Epiphytic algae were sampled by carefully removing aquatic macrophytes with a sharp razor blade and transferred into polythene bags. Distilled water was then poured into it and agitated vigorously to remove algal materials as suggested by Foester and Schlichting (1965). The algal material was then transferred into 250ml bottles and preserved in 4% unbuffered formalin solution prior to identification. Epilithic algae were sampled following the method suggested by Douglas (1958). In this method, stones surfaces from the bank of the river were measured with a ruler and algal materials scrapped using razor blade. These were collected, transferred into sampling bottles. Distilled water and drops of 4% formalin were added and agitated to preserve it for identification.




Figure: 1. Map of Shadawanka Military Barracks Showing Shadawanka River

Analysis of Samples

Physicochemical attributes such as temperature, pH and transparency were measured *in situ* using an ordinary mercury bulb thermometer, a portable Cyber Scan pH meter, model pH²⁰ and a 12cm diameter Secchi disc respectively. Electrical conductivity using a combo conductivity meter model, h1-98129, total dissolved solids by evaporation method as described by APHA (1985). Dissolved oxygen titrimetrically Using Wrinkler's method, Nutrients were determined Spectrophotometrically using an Atomic Absorption Spectrophotometer Model VGP 210 and Biochemical Oxygen Demand by Hunts Method.

Results

Biological Indices

a. Species Diversity

This was employed to determine the algal community structure in the river. The measure of species richness was given by Antoine *et al.*, (1996) as:

$$R = \frac{S}{\sqrt{N}}$$

Where; S = Number of species
N = Total number of individuals in the sample

b. Coefficient of Similarity between Algal Communities

The coefficient of similarity of the three algal communities namely; Phytoplankton, Epiphyton and Epilithon were determined using the Index of Similarity (S) between each two samples, given as;

$$S = \frac{2C}{A+B} \quad (\text{Odum, 1971})$$

Where;

A = total number of algae in community A

B = total number of algae in community B

C = number of algae common to Community A and Community B

Species Composition and Relative Abundance of Algal Taxa

A total of 756 algal species (1) were identified belonging to six classes namely; Bacillariophyceae (Diatoms), 431 species (57.0%), which formed the bulk of the algal species; the Chlorophyceae, 218 species (27.0%), Cyanophyceae, 57 species (7.5%), Euglenophyceae, 54 species (7.1%), Xanthophyceae, 7 species (1.0%) and Phaeophyceae, 3 species (0.4%).

Species Richness ("R") of algae in Shadawanka River

The Species Diversity as represented by Species Richness 'R' is presented in Table 1. The highest species richness (16.50) was obtained in the phytoplankton and the lowest (3.82) was in the Epilithon whilst Epiphyton was in between (10.51).

Table 1. Species Richness ('R') of Algal Taxa

Algal Communities	Species Richness ('R')
Phytoplankton	16.50
Epiphyton	10.51
Epilithon	3.82

The Coefficient of Degree of Similarity ("Cz") of algae in Shadawanka River

The Coefficient of Degree of Similarity ("Cz") between the algal communities in the river is presented in Table 2. The similarity between phytoplankton and Epiphyton and that between Epiphyton and Epilithon were same (0.17), whilst the similarity obtained between the phytoplankton and Epilithon communities was lower (0.14).

Table 2. Coefficient of Degree of Similarity between Algal Communities

Algal Communities	Coefficient of Similarity
Phytoplankton Vs Epiphyton	0.17
Phytoplankton Vs Epilithon	0.14
Epiphyton VS Epilithon	0.17

Table 3 Summary of the Physicochemical Parameters of Shadawanka River

Parameter	Station A			Station B			Station C			Station D		
	Mean ± S.E	Min	Max	Mean ± S.E	Min	Max	Mean ± S.E	Min	Max	Mean ± S.E	Min	Max
Temp(°C)	24.2±3.59	20.2	28.6	23.7±2.87	20.6	27.2	24.5±2.09	20	27.3	23.6±3.34	20	28.1

Transp(m)	0.22±0.15	0.11	0.3 3	0.36±0.12	0.22	0.64	0.33±0.06	0.2 0	0.4 3	0.40±0.09	0.2 5	0.5 5
pH	7.06±0.15	6.8	7.3 0	6.90±0.45	6.1	7.3	7.11±0.28	6.9	7.7	7.05±0.27	6.5	7.5
NO₃- N(mg/L)	6.65±0.52	6.01	7.1 5	7.13±0.68	6.33	8.0	6.71±0.37	6.2 4	7.0 2	6.42±0.38	6.0	6.8 8
PO₄- P(mg/L)	0.32±0.06	0.22	0.4 4	0.26±0.06	0.22	0.53	0.20±0.07	0.1	0.2 8	0.23±0.08	0.1	0.3 2
DO(Mg/L)	12.73±2.12	10.0	15. 8	12.18±1.5 7	10.2	15.9	11.96±1.35	9.8	13. 4	10.97±0.87	9.8	11. 9
TDS(mg/L)	0.85±0.24	0.5	1.2	2.86±4.32	0.3	14.3	3.02±4.16	1.0	12. 1	3.35±4.23	0.2	11. 7
BOD(mg/ L)	4.79±0.97	4.8	6.4	4.68±1.17	3.1	6.0	4.38±0.42	3.9	4.2	3.84±0.42	2.1	5.0
EC(µmSc⁻¹)	450.6±211.4	210	920	449.6±192	230	780	412.6±82.8	204	580	415.2±92.5	245	840
	6		.6				3			6		

CONCLUSION

Generally, the most marked trend of the wet season over the dry season was; increased Conductivity, Depth, and Phosphate, then decrease in Temperature and Transparency. Other water quality parameters such as Nutrients showed no pronounced temporal trends.

Temperature varied significantly with stations. Station A and D recorded the highest temperature among d stations. Station C recorded the least in the dry season while station A and C had the highest temperature in the dry season which was probably due to changes in depth and time of sampling. Temperatures were virtually same in September but generally lower temperatures fall within the months of July at stations B, C, and D. Station A in August. Lower transparency was observed at station C in June, it varied greatly among all the stations. Transparency was generally low among stations B, C, and D, in the months of April to September. The highest transparency recorded was at station A. There was no definite pattern among the stations, this could be due to time it was determined. pH is invariably same among all the stations. As a general trend, there were no much variations from acidity to alkalinity in station A, and B. A vivid variation was observed in September where acidity was maintained at stations C and D probably due to deposits of waste and domestic inputs like detergents and fertilizers. Nitrate and phosphate varied greatly among all the stations, this could be as a result of increase and decrease in levels of eutrophication. Dissolved Oxygen however, did not vary much among the stations, it had almost same trend of change of change within each station throughout the period of study. Total dissolved Solids had a great variation through both seasons and among stations with the greatest variation at station D in the dry season and station B in the wet season. Electrical Conductivity varied greatly among all the stations, it increased with the wet season. Biochemical Oxygen Demand however did not vary significantly among the stations in both seasons.

Higher numbers of algal biomass were observed towards the end of the year (September) and towards the end of the dry season (March). Bacillariophyceae dominated other algal taxa on several occasions by the

pinnate and centric forms. Diatoms dominated both the Epiphyton and Epilithon communities too in the total algal biomass, however, differences were noted during the wet season where less or greater contributions to the biomass of other algal taxa is seen. This trend was observed by Antoine (1995), Ezra (1995) and Kadiri (1993) on the existence of more than one period of diatom activity.

Chlorophyceae, the green algae came second in abundance to Bacillariophyceae, there was an increase in diatoms and desmids though, fewer numbers were found in the wet season, in varying proportions among the algal communities, these could be as a result of increased flow and high turbidity (low transparency) as reported by Antoine (1995) and Watanabe (1977). The occurrence of higher number of desmids is an attestation of low nutrient status of the river since they are a characteristic of freshwater environment with poor ionic composition as reported by Kadiri (1993 and 1999).

Occurrence of Cyanophyceae was barely inconsistent during the period of study in both seasons which were common to Phytoplanktonic communities only. It could be possible that, the river is regulated by the flow from station A and its pH values favoured the growth of this group in the river other than on substrate preferences. Cyanophyceae in small number are a natural part of the water system. In large number, they spoil the water system (Thomas, 2000).

Euglenophyceae had almost the same pattern of abundance. Euglenophyceae and Cyanophyceae also had almost the same pattern of abundance with each other and were quantitatively less numerous (7.1% and 7.5% respectively). These two groups are characteristics of eutrophic or nutrient rich bodies; their poor representation therefore, is rather, a corroboration of poor nutrient status of the river as reported by Kadiri (1996). Phaeophyceae and Xanthophyceae were sparsely represented during the study period.

Most increase in the algal population coincided with periods of longer sunshine hours of increased sunlight intensity and water temperature with low flow and slow current of the river. However, light intensity does not seem to play a major role, since higher intensity reached the river bed during the dry season due to high transparency. This was also reported by Antoine *et al.*, (1996).

Temperature, pH, phosphate and electrical conductivity were highly insignificant when correlated with the total algal cells; this indicates that, they were not limiting factors or playing any major role in limiting the algal growth, however, Transparency, Nitrate, Dissolved Oxygen, Depth and Biochemical Oxygen Demand, Total Dissolved Solids variations showed a significant correlation ($P < 0.05$) with algal variables. Physical characteristics and rainfall may be the main contributions factors, which affected the algal growth

Species richness was high probably indicating the contribution of many species in the total algal biomass. According to Nwankwo (1996), low values of diversity index may indicate dominance by one or two species: and high value, when the numbers of species are more evenly distributed (Dispersed) the relatively low Species Richness in Epilithon could be as a result of few rocks sampled. The Coefficient of Degree of Similarity between Phytoplankton and the Epiphytic communities might have resulted from the frequent attachment of algae to the macrophytes and the dislodgement due to wind activities and water turbulence. Similar findings were observed by Ezra (1995) and Nwankwo (1998).

REFERENCES

- Antoine, EA. (1995). Epilithic Algal Population of the River Wye System, Wales, U.K. *Journal of Algological Studies* 78:11-31.

- Antoine, EA. Bowker, D. W and Anyam, R. W. (1996). The Relationship between the Suspended and Periphytic Algae of River Ithon Wales UK *Limnologia* 20 (1) 73-82
- American Public Health Agency (1985). Standard Methods for the Examination of Water and Waste Water, 15th Edition, American Public Health Association, Washington, D.C pp: 1134.
- Ezra, AG. (1995). The Ecological Studies of Algae in Gubi Dam, Bauchi State, Nigeria...PhD Thesis, Abubakar Tafawa Balewa University Bauchi, Bauchi State, Nigeria p. 187
- Kadiri, MO. (1993). Further Desmids from Ikpoba Reservoir (Nigeria) Compared with other Records from. Africa-*Algological Studies*.71:23-35.
- Kadiri, MO. (1996). More Desmids from Ikpoba Reservoir Nigeria Compares with other African Records, *Algological Studies* 71:87-98.
- Kadiri, MO. (1999). A Comprehensive Preliminary Checklist of the Algae of the Ikpoba Reservoir, Edo State *Global Journal of Pure and Applied Sciences* 5:485-491.
- Moss, B. (1973). The Influence of Environmental Factors of Freshwater Algae, An Experimental Study III Effect of Temperature, Vit, Requirements and Organic N Compound on Growth *Journal of Ecology* 61:179-192.
- Nwankwo, DI (1990). Contributions to Diatom Flora of Nigeria (1) Diatoms of Lagos Lagoon and Adjacent Sea *Nig. Journal of Botany* 3:187-207.
- Nwankwo, DI (1998). Seasonal Changes in Phytoplankton Composition and Diversity, *Arch, Hydrobiologia* 40 (2) 83-93.
- Odum, E. P. (1971) Fundamentals of Ecology: W. B. Saunders Company, Philadelphia, USA pp.574
- Patrick R. (1988). Importance of Diversity in the Functioning and Structure of Riverine Communities *Limnol. Oceanogr* 33:1304-1307.
- Reynold, C. S (1984). The Ecology of Freshwater Phytoplankton, Cambridge University Press p.384.
- Thomas, M. (2000). Harmful Algal Bloom in North Carolina Clear Creek Farm, <http://www.epistate.nc.us/epi/lab/blueGreen.html> Occupational and environmental Epidermology.
- Watanabe, T. (1977). Water Pollution of Kanzaki-Gara River bin Osaka Prefecture and Diatom Flora of the Bottom Mud on the River Bed. *Nova. Hydrobiologia*. 6:27-65.
- Watson, S.B, Cauley, M.C, and Downing, J.A (2007). Pattern in Phytoplankton Taxonomic Composition across Temperate Lakes of Differing Nutrient Status *Limnol* 36:64-78.