

Int. J. Forest, Soil and Erosion, 2016 6 (1)

ISSN 2251-6387

© February 2016, GHB's Journals, IJFSE, Shabestar, Iran

Research Paper

### Investigation on the effects of *Salsola arbusculiformis* Drob. on soil properties in North East rangelands of Iran

Ali mohammad asaadi<sup>\*1</sup>, Gholam ali heshmati<sup>2</sup> and Ali reza dadkha<sup>3</sup>

\*-1Ph.D. Student of Range and Watershed management Faculty, Agriculture and Natural Resource University of Gorgan, Iran. E-mail: am-asaadi@um.ac.ir

2 Prof., Range and Watershed management Faculty, Agriculture and Natural Resource University of Gorgan, Iran. E-mail: heshmati.a@gmail.com

3Associate Professor, High Education center of Shirvan. E-mail: Dadkha@um.ac.ir

**Abstract:** The purpose of this investigation was to study the effects of *Salsola arbusculiformis* on soil physico-chemical characteristics in rangelands of Garmeh Located in northeast of Iran. After surveying the regions, the three sites (Karnakh, Aspakho, Robat) with *Salsola arbusculiformis* were selected. Vegetation cover and soil parameters were sampled by a random systematic method, in each region. Five transects each 100 meter length were selected and then 50 plots by the average of 1m × 2m were randomly chosen and vegetation cover parameters such as percentage canopy cover, production, density and percentage litter were measured in the key area of each site. Soil samples were taken from two depths (0-30, 30-60 cm) under plant and in control areas. The measured characteristics of soil included texture, electrical conductivity (EC), acidity (pH), nitrogen (N), phosphorus (P), potassium (K), Sodium (Na), Organic matter (OM), CaCO<sub>3</sub> and CaSO<sub>4</sub>. Data were analyzed using SPSS software under windows. One-way ANOVA and T-test were used to compare the data related to planted and control treatments. Results showed that *Salsola arbusculiformis* significantly increased the amount of K, pH, EC, N, Na (P<0.01), and O.M (P<0.05). In these sites measured characteristics of soil in planted areas were more than the control ones.

**Key words:** *Salsola arbusculiformis*, Soil physico-chemical properties, Garmeh, Iran.

#### Introduction

Knowledge of ecological factors such as climate, soil, topography and disturbance influencing plant species distribution is essential for conservation, management, and recovery of rangeland ecosystems (Ajeer & Shahmoradi, 2007; Nautiyal et al., 2009). Generally ecosystem is a unit consisting of all factors, including environmental factors and organisms in which the interaction of animate and inanimate components should be considered. However, the vast number of rangeland ecosystems and the variety of variables brings out estimating complexities of relationships and their effects. Therefore, effective relationships between plants and the environmental context can be studied under different categories. One of these items is the effect of plants on soil. Plants can be indicators of ecological conditions that show a real ecological in each region (Khedri Gharibvand, 2007). The interaction of soil-plant and morphological and physiological status of plants determines the dominant effect of plant or soil. Therefore, such environments combined together in a state of steady flow are placed in the way of activities, actions and interactions which affect and influence each other to create the change. According to studies conducted, the establishment of vegetation and its expansion causes the improvement of soil physical and chemical characteristics and facilitate natural resource management particularly rangelands. Some of environmental factors such as moisture and soil nutrients had significant effect on plant-community (Mahdavi Ardakani et al., 2010).

Plants affect physical and chemical characteristics of soil. It can be said that the food reserves in the soil properties and vegetation is highly dependent on plant species (Barth, 1980). The soil characteristics are affected by soil response to root activity and characteristics of litter that fall under canopy collapses perennial plants (Balamurgan et al., 2000). For example, perennial plants by accumulation of litter and whose roots affect the soil quality, improve their sites (Davidson and Morton, 1984). The effect of plants on soil characteristics has been studied in Iran and other parts of the world.

Bailey (1970) believes the vegetation, will speed the nutrient cycle and created favorable changes in the microclimate of the area. Mugunga and Mugumo (2013) studied the effects of *Acacia sieberiana* plants on the soil chemical properties in Songa, they reported that the amounts of nitrogen, Calcium, soil organic carbon, Cation exchange capacity (CEC) and pH were higher in soils under tree canopies than open areas. Halvarson et al. (1997) in the study of sagebrush fields south of Washington concluded that the accumulation of organic carbon and nitrogen and the speed of nitrogen cycle in the soil below of *Artemisia tridentata* plant is more than the soil between plants.

Jafari et al., (2006) in rangelands of Qom province, Iran, studied the relationships between soil properties and plant species distribution. Their results showed that the most important factors for separation of plant types were soil texture, electrical conductivity and CaCO<sub>3</sub>.

Mojiri and Jalalian (2011) results that electrical conductivity, sodium ions, HCO<sub>3</sub><sup>-</sup> and the percentage of clay had negative effects on *Nitraria schoberi* plant physiologic parameters. Koochaki (1996) mentioned some reasons for differences in soil nutrient levels under the crown of the plant as follow: 1) nutrient uptake by plant roots shallow and deep, 2) immobilization of nutrients by plants or symbiotic organisms, 3) increased soil nutrients by large organisms that use plants to their nests and rest, etc and 4) stop and accumulate in Brushwood and soil particles caused by wind at the foot of shrubs and trees. Jahanbin et al. (2013) reported that *Myrtus communis* plants increased organic matter, total nitrogen, potassium, phosphorous, copper, manganese, iron, zinc and electrical conductivity of soil. The purpose of this investigation was to study the effects of *Salsola arbusculiformis* on physico-chemical characteristics of soil in Robate, Aspakho and Karnakh regions in North East rangelands of Iran.

## MATERIALS AND METHODS

### Study area:

The study area was chosen in arid rangelands of Garmeh in North East of Iran (North Khorasan Provinces). This region is situated between 37° 18' to 37° 24' North latitude and 56° 20' to 56° 37' East longitudes. The area is approximately 30000 hectares with elevation ranging from 1280 m to 1600 meter. The means of precipitation is 229mm/year that maximum and minimum of precipitation occur in April and July respectively. The mean of annual temperature is 12.9 °C. The average maximum temperature is 32.6 °C in July and minimum temperature is -5.92 °C in January. The climate of this region with using of Emberger method is cold arid. The embrothermic diagram show that drought period is for six months of year and wet season start in November and continues until May.

After identifying the regions, the three sites (karnakh, Aspakho, Robat) with *Salsola arbusculiformis* were selected. Five transects each 100 meter length were selected and then 50 plots by the average of 1m ×2m were randomly chosen and vegetation cover parameters such as percentage canopy cover, production, density and percentage litter were measured in the key area of each site. Soil samples were taken from two depths (0-30, 30-60 cm) under plant and in control areas. Soil samples after drying and passing the 2 mm sieve to determine chemical properties included texture, electrical conductivity (EC), acidity (pH), nitrogen (N), phosphorus (P), potassium (K), Sodium (Na), Organic Carbon(OC), CaCO<sub>3</sub> and CaSO<sub>4</sub>. Data were analyzed using SPSS software under windows. One-way ANOVA and T-test were used to compare the data related to planted and control treatments.

### RESULTS AND DISCUSSION

Climate is one of the most important factors in the plant development (Vallejo et al., 1998). Irano-Turanian zone which fall and winters are cold, rainy and snowy, and summers are dry and hot, are covered by sclerophylleous species that need quite little water and high temperatures (Dogan et al., 2003). Using the Emberger climate classification, the habitat of *Salsola arbusculiformis* was categorized as cold arid. The mean annual rainfall and temperature were about 229 mm and 12.9 °C for the habitat, respectively, with the bulk of the rains concentrated in February, March, April and May.

The Flora of study area is rich composed of annual and perennial plants. This area is categorized as fair rangelands. Other plants include *Artemisia sieberi*, *Poa bulbosa*, *Stipa barbata*, *Carex sp.*, *Eurotia ceratoides*, *Salsola rigida*, *Bromus tectorum* and *Scariola orientalis*.

**Soil texture:** The soil particles size composition of soil in the area is shown in Table 1. The physical analysis of the soils showed that *Salsola arbusculiformis* grows on loamy (63.34%), sandy-loam (18.33%), clayey-loam (15%) and Silt-loam (3.33%) soils. The results showed that the soil texture under the *Salsola* canopy hasn't changed. Soil texture has a large effect on controlling soil moisture and nutrients available to plants (Jafari et al., 2006). Among morphological parameters, soil and vegetation are closely related. Therefore, any suggestion for reform in rangeland should take into account the soil properties. Having the knowledge of soil characteristics of each species has effective role in suggesting compatible species to the soil conditions in the same areas (Jafari et al., 2006).

**Acidity (pH):** The results indicated that among of Acidity in the studied sites are significantly different (Table1). In relation to the effects of *Salsola arbusculiformis* on soil acidity results showed that soil pH is significantly increased under *Salsola* canopy rather than outside it and save it under 8.93 and outside the canopy 8.15 units (Table 2). Soil analysis show that soil samples are moderately alkaline, according to JACKSON's (1958) soil-pH classification. The pH level of the soil surface was significantly higher than the soil depth. There was an interaction between soil depth and soil cover dressing so that with increasing the depth, shading acidity increased (Table 2). Zheng et al. (2008) while studying *Salsola passerina* found that the pH with increasing depth plants did not show significant differences, but its rate savings under the canopy is 9.1 and outside the 8.45.

**Electrical conductivity (EC):** *Salsola arbusculiformis* significantly increased the electrical conductivity. EC under *Salsola* canopy was 590 versus 200 μS/cm out of *Salsola* canopy. EC indicate significant decreases with increasing depth (Table 2). In comparison, the interaction of different levels of electrical conductivity depth coverage on the outside of the canopy, there is no significant difference, but under different soil depths in contrast to significant savings can be seen that the electrical conductivity decreased with increasing depth (Table 1). Plants existences provide shade and reduce the temperature of the soil surface evaporation, less water and less solute transfer from the surface to depth, while the organic acids resulting from the decomposition of organic material dissolved minerals and helps release ions (Titus et al., 2002). Falah Shojaee (2006) in the study of four *Acacia* species concluded that the electrical conductivity of soil in *Acacia* canopy shading were significantly higher than out plants and they express that most of activity of micro-organism in shading due to higher food region, leading to increased secretion of organic acids in the root zone, which is an increase of ions causes an increase in electrical conductivity. Results Everett et al. (1986) showed that the electrical conductivity of the soil-plant pine trees out there and also shading and gradual decrease of the electrical conductivity decreased with depth. McDaniel and Graham (1992) indicate that electrical conductivity beneath the canopy and outside canopy of pine has a significant positive correlation with increasing depth. Zheng et al. (2008) found that the rate of *Salsola passerina* rangeland plant canopy under increased electrical conductivity and the electrical conductivity decreases dramatically with increasing depth.

**Organic carbon:** The results showed that among of Organic carbon in the studied habitats are significantly different (Table1). *Salsola arbusculiformis* significantly increased soil organic carbon under the plant canopy (Table 2). So that out of the shadows is 0.79% and savings under 1.03%. Organic carbon in the soil surface is higher than the lower layers. Mlambo et al. (2005) in South Africa to study *Colophosprum mopane* plant amount of organic carbon in the canopy obtained from zero to 10 cm depth was significantly higher than that obtained from shading. The increase in organic carbon can be attributed to various processes such as accumulation of litter (Zinke, 1962), reducing in erosion or increase in sedimentation (Charley, 1977), improve micro-climatic soil conditions (Pierson and Wight, 1991) or be input sources such as insects, birds and other animals (Davenport et al., 1996). In general, the amount of organic matter accumulated Litter plant size and there is a direct relationship (Titus et al., 2002).

**Nitrogen (N):** *Salsola arbusculiformis* has a significant impact on canopy soil nitrogen. As can be seen in Table 2, the amount of soil nitrogen was higher under *Salsola* canopy. Nitrogen levels in the canopy soil and the outside canopy soil was 0.03 and 0.025 %. This difference significantly different between the depth can be seen in the surface soil nitrogen grenades in the verse above (Table 1).

The contrasts between the different depths and soil cover have difference significant, and the effect on Common Myrtle plant phosphorus in the soil indicates (Table 2).

Table 1: The analysis results of soil in the habitats of *Salsola arbusculiformis*.

Soil properties	Region							
	Soil depth (cm)	Robate		Aspakho		Karnakh		
		under canopy	outside canopy	under canopy	outside canopy	under canopy	outside canopy	
EC ( $\mu\text{S}/\text{cm}$ )	0-30	546a	177b	698a	206b	522a	<b>213b</b>	
	30-60	549a	305b	420a	179b	509a	<b>157b</b>	
PH	0-30	8.764b	8.19c	9.15a	8.122c	8.864b	<b>8.134c</b>	
	30-60	8.204b	8.072b	8.784a	8.128b	8.6ac	<b>8.23cb</b>	
saturation moisture %	0-30	52.968a	45.21a	52.418a	47.71a	44.712a	<b>45.966a</b>	
	30-60	59.046a	49.442a	51.72a	51.682a	48.712a	<b>43.61a</b>	
CaCO <sub>3</sub> %	0-30	25.37a	21.92a	26.385a	24.25a	24.13a	<b>23.075a</b>	
	30-60	22.845a	22.795a	24.27a	25.305a	21.587a	<b>25.505a</b>	
CaSO <sub>4</sub> %	0-30	0.886a	0.878a	0.883a	0.879a	0.887a	<b>0.884a</b>	
	30-60	0.893a	0.879a	0.886a	0.878a	0.892a	<b>0.880a</b>	
OC%	0-30	1.33a	1.13ab	1.37a	1.19a	0.94ab	<b>0.66b</b>	
	30-60	0.94a	0.697a	1.04a	0.631a	0.53a	<b>0.482a</b>	
N%	0-30	0.0284a	0.0293a	0.0297a	0.0241bc	0.0255b	<b>0.0227c</b>	
	30-60	0.0242a	0.0239a	0.0227a	0.0205bc	0.0216ab	<b>0.0194c</b>	
K(mg/kg)	0-30	81.643a	40.46d	101.37b	40.155d	58.903c	<b>40.884d</b>	
	30-60	31.656bc	20b	65.53a	25.093b	41.468c	<b>28.42bc</b>	
P(mg/kg)	0-30	0.052a	0.123a	0.079a	0.78a	0.136a	<b>0.171a</b>	
	30-60	0.064a	0.074a	0.06a	0.082a	0.05a	<b>0.03a</b>	
Na(mg/kg)	0-30	67.02a	4.74b	52.95ac	5.63b	40.16c	<b>8.40b</b>	
	30-60	76.30a	20.52b	73.95a	5.92b	52.50a	<b>4.23b</b>	
Clay (%)	0-30	10.35	10.28	10.47	13.14	12.38	<b>14.31</b>	
	30-60	25.74	23.65	26.47	19.37	23.65	<b>24.04</b>	
Silt (%)	0-30	38.93	41	30.81	37.74	39	<b>31.6</b>	
	30-60	37.94	36.03	40.01	34.71	41.73	<b>37.84</b>	
Sand (%)	0-30	50.72	48.72	58.72	49.12	48.62	<b>54.12</b>	
	30-60	36.32	40.32	33.52	45.92	34.62	<b>38.12</b>	

Note: Values with unlike letter in a row are significantly different at 0.05 levels.

**Potassium (K):** The results showed that among of Potassium in the studied habitats are significantly different (Table1). In this study, soil K under *Salsola* canopy was significantly higher (80.64 mg kg<sup>-1</sup>) than outside the canopy (40.5mg per kg). This increase can also be seen at various depths in the soil (Table 2). Mishra et al. (2003) showed a significant difference in soil K between the canopy and outside canopy of 3 and 6 years old eucalyptus trees. The reason for this increasing could be due to releasing potassium from K-bearing minerals and causes significant increase in K under the canopy of K release or litter decomposition relationships. Banerjee (1986) with the ecosystem studying of which it was the dominant oak species, expressed as the amount of exchangeable potassium in the canopy than outside the shading. Karimian and Razmi, (1990) causes an increase in the concentration of potassium in the plant canopy organic matter in the plant shading savings and increased biochemical activity and results in the release of potassium-bearing minerals potassium. Wang et al. (2000) described an increase of organic acids secreted by plant roots leads to release of potassium by plants contain minerals such as potassium feldspar gneiss and is. Tan (1978) effect of organic compounds on Humic acid and Folic acid and K release from illite and smectite clays showed.

**Phosphorus (P):** There was no significant difference ( $P>0.05$ ) in the soil Phosphorus between the control and under the canopy soil. However, Phosphorus ranges were between 0.02 and 0.09 mg/kg. The average value for soil P is between 0.0006 and 0.0009% (Eskin et al., 2013). The results showed that soil P for three locations is low.

**Sodium (Na):** The results indicated that among of Na in the studied sites are significantly different (Table1). *Salsola arbusculiformis* significantly increased the amount of sodium ions under the plant canopy than outside the canopy (Table 2). Soil sodium increases soil reaction, encourages low uptake in some types of micronutrients and destroys the physical properties of soils, thereby reducing soil permeability and growth of plants (Honarjoo et al., 2010).

**The Percentage of Calcium Carbonate (CaCO<sub>3</sub>):** The soils habitats of this plant have high level lime varying between 14.5 to 29.15%. According to Table 2, the percentage of calcium carbonate was no significant difference ( $P>0.05$ ) between the control and under the canopy. Soil carbonate causes the creation of appropriate soil structure and induces changes to soil acidity (Jafari et al., 2006).

**The Percentage of CaSO<sub>4</sub>:** There was no significant difference ( $P>0.05$ ) in the soil CaSO<sub>4</sub> between the control and under the canopy soil. The soils habitats of this plant have very low level gypsum (Table1).

**Saturation moisture percentage:** Tests showed that the *Salsola arbusculiformis* shrubs increased saturation percentage. Saturation percentage below perspective shadows is 51.59% and out of perspective shadows is 47.26%. But it is not a significant difference between different regions and soil depths (Table 1, 2).

**CONCLUSION**

The results show that the *Salsola arbusculiformis* can influence soil chemical and physical characteristics of their habitat. In this study, the shrubs increased electrical conductivity, acidity, organic carbon, potassium, sodium, percent nitrogen and saturation. It can be impressively used for rehabilitation of desertified rangeland and improvement of degraded rangeland in arid regions. In addition, *Salsola arbusculiformis* could be a good fodder resource for sheep and goat. Six month of vegetative growth in this species is grazing time.

Table 2: The compare of physical and chemical analysis results on the soil in the habitats of *Salsola arbusculiformis* in 0-30 and 30-60 cm depths.

Soil properties	Treatment	Average 0-30 (cm) depth	Average 30-60 (cm) depth	SE	sig.
EC ( $\mu\text{S}/\text{cm}$ )	under canopy	590	490	66	ns
	outside canopy	200	210	35	ns
PH	under canopy	8.93	8.53	0.125	**
	outside canopy	8.15	8.14	0.038	ns
saturation moisture %	under canopy	50.03	53.16	2.09	ns
	outside canopy	46.29	48.24	2.38	ns
CaCO <sub>3</sub> %	under canopy	25.29	22.9	1.052	ns
	outside canopy	23.08	24.54	1.02	ns
CaSO <sub>4</sub> %	under canopy	0.88	0.89	0.0038	ns
	outside canopy	0.88	0.88	0.0012	ns
OC%	under canopy	1.21	0.84	0.15	*
	outside canopy	0.99	0.6	0.10	**
N%	under canopy	0.03	0.023	0.0006	**
	outside canopy	0.025	0.021	0.0004	**
K(mg/kg)	under canopy	80.64	46.22	6.48	**
	outside canopy	40.5	24.5	1.67	**
P(mg/kg)	under canopy	0.09	0.06	0.022	ns
	outside canopy	0.12	0.06	0.019	ns
Na(mg/kg)	under canopy	53.37	67.58	8.93	ns
	outside canopy	6.25	10.22	2.78	ns
Clay (%)	under canopy	11.07	25.38	1.85	**
	outside canopy	12.57	22.35	1.77	**
Silt (%)	under canopy	36.25	39.89	2.08	ns
	outside canopy	36.78	36.19	3.02	ns
Sand (%)	under canopy	52.7	34.82	3.07	**
	outside canopy	50.65	41.45	3.28	*

\*Significant at 5% alpha level, \*\*Significant at 1% alpha level

**Reference**

- Ajeer F, Shahmoradi A (2007). Autecology of Range plants species of *Ferula ovina* in Tehran Province. Iranian Journal of Range and Desert Research. 14(3): 367-359.
- Bailey A W (1970). Barrier effect of the shrub *Elaeagnus commutate* on grazing cattle and forage production central Alberta. Journal of Range management. 23(4): 248- 251.
- Balamurgan J, Kumaraswamy K, and Rajarjan A (2000). Effects of *Eucalyptus citriodora* on the physical and chemical properties of soil. Journal of the Indian Society of Soil Science. 48(3): 491-495.
- Barth R C (1980). Influence of pinyon pine trees on soil chemical and physical properties. Soil Science Society of America Journal. 44: 112-114.
- Banerjee S K, Nath S, Banerjee S P (1986). Characterization of soils under different vegetations in the Tarai region of Kurseong forest division, West Bengal. Journal of the Indian Society of Soil Science, 34(1): 343-349.
- Charley J L, West N E (1977). Micro-patterns of nitrogen mineralization activity in soils of some shrub-dominated semi-desert ecosystems of Utah. Soil Biology and Biochemistry. 9:357-365.
- Davenport D W, Bradford PW, Breshears D D (1996). Soil morphology of canopy and intercanopy sites in a pinon-juniper woodland. Soil Science Society of America Journal.60: 1881-1887.
- Davidson D W, Morton S R (1984). Dispersal Adaptations of Some Acacia Species in the Australian Arid Zone. Ecology. 65:1038-1051.
- DoganY, Baslar S, Aydin H, Mert H (2003). A study of the soil-plant interactions of *Pistacia lentiscus* L. distributed in the western Anatolian part of Turkey. Acta Botanica Croat. 62: 73-88.
- Eskin B, Ozyigit I, Dogan I, AltayV, Demir G, Serin M (2013). Germination Physiology and Autecology of *Centaurea kilaia* Boiss. from Turkey. Sains Malaysiana. 42(10): 1473-1482.
- Everett R L, Sharrow S H, Thran D (1986). Soil nutrient distribution under and adjacent to single leaf crowns. Soil Science Society of America Journal. 50:788-792.
- Falah Shojaee J (2006). Effect of Acacia species on soil physical and chemical properties of plain corruption Grbaygan city. Master's thesis, Soil Science. Faculty of Agriculture. University of Shiraz.

- Halvarson J, Bolton H, Smith J (1997). The pattern of soil variables related to *Artemisia tridentata* in burned shrub-steppe site. Soil Science Society of American Journal. 61: 287-294.
- Honarjoo N, Mojiri A, Jalalian A, Karimzadeh H R (2010). The effects of salinity and alkalinity of soil on growth of *Haloxylon sp.* In Segzi plain (Iran). International Conference on Chemistry and Chemical Engineering (ICCCE 2010), Kyoto, Japan 1-3 August: 285-288.
- Jackson ML (1958). Soil Chemical Analysis. New Jersey: Prentice Hall.
- Jafari M, Zare Chahuki M A, Azarnivand H, Baghestani N, Tavili A, Kohandel A (2006). The relationship between soil features and plant species in rangeland of Qom province. Research and Development. 73:110-116.
- Jahanbin R, Jahantab E, Alirezanezhad A, Javdani Z, Mirzaee M (2013). The effects of shrubs common myrtle (*Myrtus communis*) on soil chemical and physical characteristics of Basht area. Annals of Biological Research. 4 (5):158-164.
- Karimian, N, Razmi K (1990). Influence of perennial plants on chemical properties of arid and calcareous soils in Iran. Soil Science 150: 717-721.
- Khedri Gharibvand H (2007). Survey on some environmental factors on distribution of *Camphorosma monspeliaca* in Chaharmahal-Bakhtiari province, M.Sc Thesis, Tehran, Tarbiat Modarres University.
- Koochaki A (1996). Plant Operation Ranch Fields (translated). Press Ferdowsi University of Mashhad.
- Mahdavi Ardakani S R, Jafari M, Zargham N A, Zare Chahuki M A, Baghestani Meibodi N, Tavili A (2009). Investigation on the effects of *Haloxylon aphyllum*, *Seidlitz rosmarinus* and *Tamarix aphylla* on soil properties in ChahAfzal-Kavir (Yazd). Iranian Journal of Forest. 2(4): 357-365.
- McDaniel P A, Graham R C (1992). Organic carbon distributions in shallow soils under pinyon-juniper. Soil Science Society of American Journal. 56:499-504.
- Mishra A, Sharma S D, Khan G H (2003). Improvement in physical and chemical properties of sodic soil by 3, 6, and 9 years old plantation of *Eucalyptus tereticornis* Biorejuvenation of sodic soil. Forest Ecology Management. 184:115-124.
- Mlambo D, Nyathi P, Mapaure I (2005). Influence of *Colophospermum mopane* on surface soil properties and understorey vegetation in a southern African savanna; Forest Ecology Management. 212: 394-404.
- Mojiri A, Jalalian A (2011). Relationship between Growth of *Nitraria schoberi* and some soil properties. The Journal of Animal & Plant Sciences. 21(2): 246-250.
- Mugunga C P, Mugumo D T (2013). *Acacia sieberiana* Effects on Soil Properties and Plant Diversity in Songa Pastures, Rwanda. International Journal of Biodiversity. 1-11.
- Nautiyal B P, Nautiyal M C, Khanduri V P, Rawat N (2009). Floral biology of *Aconitum heterophyllum* Wall: A critically endangered alpine medicinal plant of Himalaya, India. Turkish Journal of Botany. 33: 13-20
- Pierson F B, Wight J R (1991). Variability of near-surface soil temperature on sagebrush rangeland. Journal of Range Management. 44:491-497
- Titus J H, Nowak R S, Smith S D (2002). Soil resource heterogeneity in the Mojave Desert. Journal of Arid Environments. 52: 269-292.
- Tan K H (1978). Effects of humic and fulvic acids on release potassium. Geoderma. 21(1):67-74.
- Vallejo V R, Serrasolses I, Cortina J, Seva J P, Valdecantos A, Vilagrosa A (1998). Restoration strategies and actions in Mediterranean degraded lands. Project report of EC Environment and Climate Programme (ENV4-CT97-0682 REDMED, Climate and Natural Hazards), Spain.
- Wang JG, Zhang FS, Zhang XL, Cao YP (2000). Effect of plant types on release of mineral potassium from gneiss · Nutrient cycling in Agroecosystems. 56(1): 45-52
- Zheng J, He M, Li X, Chen Y, Li X, Liu L (2008). Effect of *Salsola passerina* shrub patches on the micro scale heterogeneity of soil in mountain grassland, China. Journal Arid Environment. 72(3):150-161.
- Zinke P J (1962). The pattern of influence of individual forest trees on soil properties. Ecology. 43: 130-133.